


## Product Datasheet

### TBK1 Antibody (orb1238307)

<b>Description</b>	TBK1 Antibody	 <p>Immunocytochemistry of NAK in MOLT4 cell...</p>
<b>Species/Host</b>	Rabbit	
<b>Reactivity</b>	Human, Mouse	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, ICC, WB	
<b>Immunogen</b>	NAK antibody was raised against a synthetic peptide corresponding to 17 amino acids form near the carboxy terminus of human NAK/TBK1. The immunogen is located within the last 50 amino acids of NAK.	
<b>Target</b>	TBK1	
<b>Preservatives</b>	NAK Antibody is supplied in PBS containing 0.02% sodium azide.	
<b>Form/Appearance</b>	Liquid	
<b>Concentration</b>	1 mg/ml	
<b>Storage</b>	NAK antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	
<b>Note</b>	For research use only	
<b>Application notes</b>	NAK antibody can be used for detection of NAK/TBK1 by Western blot. An 84 kDa band should be detected. Antibody can also be used for immunocytochemistry starting at 10 µg/mL. Antibody validated: Immunocytochemistry in human samples. All other applications and species not yet tested.	
<b>Isotype</b>	IgG	
<b>Clonality</b>	Polyclonal	
<b>MW</b>	84 kDa	
<b>Uniprot ID</b>	<a href="#">Q9UHD2</a>	
<b>NCBI</b>	<a href="#">NP_037386</a>	
<b>Dilution Range</b>	NAK antibody can be used for detection of NAK/TBK1 by Western blot. An 84 kDa band should be detected. Antibody can also be used for immunocytochemistry starting at 10 µg/mL. Antibody validated: Immunocytochemistry in human samples. All other applications and species not yet tested.	