

Product Datasheet

TNFSF10 Antibody (orb1238614)

Description

TNFSF10 Antibody

Species/Host

Rabbit

Reactivity

Human

Conjugation

Unconjugated

Tested

ELISA, IF, IHC-P, WB

Applications

Immunogen

TRAIL antibody was raised against a peptide corresponding to 17 amino acids near the carboxy terminus of human TRAIL. The immunogen is located within the last 50 amino acids of Trail.

Target

TNFSF10

Preservatives

Trail Antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/ml

Storage

Trail antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note

For research use only

Application notes

TRAIL antibody can be used for detection of TRAIL by Western blot at 1 - 5 µg/mL dilution. Antibody can also be used for immunohistochemistry starting at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype

IgG

Clonality

Polyclonal

MW

Predicted: 32 kDa Observed: 35 kDa

Uniprot ID


[P50591](#)

NCBI


[NP_003801](#)

Dilution Range


TRAIL antibody can be used for detection of TRAIL by Western blot at 1 - 5 µg/mL dilution. Antibody can also be used for immunohistochemistry starting at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human



Western blot analysis of TRAIL in HeLa c...



Immunohistochemistry of TRAIL in human b...



Western blot analysis of TRAIL in human ...