

Product Datasheet

TICAM2 Antibody (orb1238769)

Description

TICAM2 Antibody

Species/Host

Rabbit

Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Tested

ELISA, IF, IHC-P, WB

Applications
Immunogen

TIRP antibody was raised against a peptide corresponding to amino acids near the C-terminus of human TIRP. The immunogen is located within the last 50 amino acids of TIRP.

Target

TICAM2

Preservatives

TIRP Antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/mL

Storage

TIRP antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note

For research use only

Application notes

TIRP antibody can be used for detection of TIRP by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human, mouse, and rat samples; Immunohistochemistry in mouse samples and Immunofluorescence in mouse samples. All other applications and species not yet tested.

Isotype


IgG

Clonality


Polyclonal

Uniprot ID
Q86XR7
NCBI
AAP81748
Dilution Range


TIRP antibody can be used for detection of TIRP by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human, mouse, and rat samples; Immunohistochemistry in mouse samples



Western blot analysis of TIRP in human (...)



Immunohistochemical staining of mouse kidney tissue



Immunofluorescence of TIRP in Mouse Kidney