

## **Product Datasheet**

STK39 Antibody (orb1238872)




## www.biorbyt.com

**⋖**Stk39

Western blot analysis of Stk39 in SK-N-S...



Immunohistochemistry of Stk39 in human b...



**Immunofluorescence** of stk39 in human bra...

Descriptionnts. STK39 Antibody

Species/Host Rabbit

Reactivity Human, Mouse

Conjugation Unconjugated

Tested ELISA, IF, IHC-P, WB

**Applications** 

**Immunogen** Stk39 antibody was raised against a 19 amino acid

synthetic peptide from near the carboxy terminus of human Stk39. The immunogen is located within

amino acids 450 - 500 of Stk39.

STK39 **Target** 

**Preservatives** Stk39 Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/mL

**Storage** Stk39 antibody can be stored at 4°C for three

> months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be

exposed to prolonged high temperatures.

Note For research use only

**Application notes** Stk39 antibody can be used for detection of stk39 by

> Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 μg/mL. For immunofluorescence start at 20 μg/mL.Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype IgG

Clonality Polyclonal

**Uniprot ID Q9UEW8** 

**NCBI** NP 037365

**Dilution Range** Stk39 antibody can be used for detection of stk39 by

> Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 μg/mL. For immunofluorescence start at 20 μg/mL.Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples

Carolina < br > 27709. United States

68 TW Alexander Drive<br/>br>Research Triangle Park<br/>br>Durham, North