

## Product Datasheet

### TACR3 Antibody (orb1239362)

**Description**

TACR3 Antibody

**Species/Host**

Rabbit

**Reactivity**

Human, Mouse

**Conjugation**

Unconjugated

**Tested**

ELISA, IF, IHC-P, WB

**Applications**
**Immunogen**

NK3R antibody was raised against a 18 amino acid synthetic peptide from near the center of human NK3R. The immunogen is located within amino acids 370 - 420 of NK3R.

**Target**

TACR3

**Preservatives**

NK3R Antibody is supplied in PBS containing 0.02% sodium azide.

**Form/Appearance**

Liquid

**Concentration**

1 mg/mL

**Storage**

NK3R antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

**Note**

For research use only

**Application notes**

NK3R antibody can be used for detection of NK3R by Western blot at 0.5 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

**Isotype**

IgG

**Clonality**


Polyclonal

**MW**


Predicted: 51 kDa Observed: 51 kDa

**Uniprot ID**
[P29371](#)
**NCBI**
[P29371](#)
**Dilution Range**


NK3R antibody can be used for detection of NK3R by Western blot at 0.5 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other



Western blot analysis of NK3R in RAW264....



Immunohistochemistry of NK3R in human br...



Immunofluorescence of NK3R in Human Brai...