Product Datasheet

ICOSLG Antibody (orb1239699)

Description	ICOSLG Antibody
Species/Host	Rabbit
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	ELISA, IF, IHC, WB
Immunogen	ICOSLG antibody was raised against a peptide corresponding to 17 amino acids near the center of human ICOSLG.The immunogen is located within amino acids 150 - 200 of ICOSLG.
Target	ICOSLG
Preservatives	ICOSLG Antibody is supplied in PBS containing 0.02% sodium azide.
Form/Appearance	Liquid
Concentration	1 mg/mL
Storage	ICOSLG antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Note	For research use only
Application notes	ICOSLG antibody can be used for Western blot at 1 µg/mL. Antibody can also be used for Immunohistochemistry at 2 µg/mL. For Immunoflorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.
Isotype	IgG
Clonality	Polyclonal
MW	Predicted: 33 kDa Observed: 60 kDa
Uniprot ID	O75144
NCBI	O75144
Dilution Range	ICOSLG antibody can be used for Western blot at 1 µg/mL. Antibody can also be used for Immunohistochemistry at 2 µg/mL. For Immunoflorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples;



Western blot analysis of ICOSLG in human...



Immunohistochemistry of ICOSLG in human ...



Immunofluorescence of ICOSLG in human ly...

Biorbyt Ltd.
7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom
Email: **info@biorbyt.com** | Phone: **+44 (0) 1223 859-353** | Fax: +44 (0)1223 280 240

Biorbyt LLC.
68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709. United States
Email: **info@biorbyt.com** | Phone: **+1 (415) 906-5211** | Fax: +1 (415) 651-8558