



## **Product Datasheet**

**Dffa Antibody (orb1239701)** 



**Description**nts. Dffa Antibody

Species/Host Rabbit

**Reactivity** Mouse

**Conjugation** Unconjugated

**Tested** ELISA, IF, IHC-P, WB

**Applications** 

Immunogen ICAD antibody was raised against a 20 amino acid

peptide near the carboxy terminus of mouse ICAD.The immunogen is located within the last 50

amino acids of ICAD.

Target Dffa

**Preservatives** ICAD Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/ml

**Storage** ICAD antibody can be stored at 4°C for three months

and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed

to prolonged high temperatures.

**Note** For research use only

**Application notes** ICAD antibody can be used for detection of of ICAD by

Western blot at 1:1000 dilution. A 45 kDa band can

be detected. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 5 µg/mL.Antibody validated: Western Blot in mouse samples; Immunohistochemistry in mouse samples and Immunofluorescence in mouse samples. All other

applications and species not yet tested.

**Isotype** IgG

**Clonality** Polyclonal

MW 45 kDa

Uniprot ID 054786

NCBI 054786


**Dilution Range** ICAD antibody can be used for detection of of ICAD by

Western blot at 1:1000 dilution. A 45 kDa band can

be detected. Antibody can also be used for

immunohistochomistry starting at 5 ug/ml For


## www.biorbyt.com



Secondary: Goat antirabbit IgG HRP conj...



Immunohistochemistry of ICAD in mouse ki...



Immunofluorescence of ICAD in mouse kidn...