



Product Datasheet

Dffa Antibody (orb1239702)



www.biorbyt.com


132 k

55 k 43 k 34 k

23 k

ICAD

Immunohistochemistry of ICAD in mouse lu...



Immunofluorescence of ICAD in Mouse Lung...

Descriptionnts. **Dffa Antibody**

Species/Host Rabbit

Reactivity Mouse

Conjugation Unconjugated

ELISA, IF, IHC-P, WB **Tested Applications**

Immunogen ICAD antibody was raised against a peptide corresponding to amino acids near the amino terminus of mouse ICAD. The immunogen is located

within the first 50 amino acids of ICAD.

Target Dffa

Preservatives ICAD Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/ml

Storage ICAD antibody can be stored at 4°C for three months

> and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed

to prolonged high temperatures.

Note For research use only

Application notes ICAD antibody can be used for detection of of ICAD by

Western blot at 1 µg/mL. A 45 kDa band can be

detected. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 10 µg/mL.Antibody validated: Western Blot in mouse samples; Immunohistochemistry in mouse samples and Immunofluorescence in mouse samples. All other

applications and species not yet tested.

Isotype IgG

Clonality Polyclonal

MW 45 kDa

Uniprot ID O54786

NCBI O54786

Dilution Range ICAD antibody can be used for detection of of ICAD by

Western blot at 1 µg/mL. A 45 kDa band can be

detected. Antibody can also be used for

mmunohistochomistry starting at 2 ug/ml

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280

Biorbyt LLC.

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North Carolina < br > 27709. United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558