



Product Datasheet

STK17A Antibody (orb1239863)



Descriptionnts. STK17A Antibody

Species/Host Rabbit

Reactivity Human

Conjugation Unconjugated

ELISA, ICC, IF, WB **Tested**

Applications

Immunogen DRAK1 antibody was raised against a peptide

> corresponding to amino acids near the amino terminus of human DRAK1. The immunogen is located within the first 50 amino acids of DRAK1.

STK17A Target

Preservatives DRAK1 Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/ml

Storage DRAK1 antibody can be stored at 4°C for three

> months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be

exposed to prolonged high temperatures.

Note For research use only

Application notes DRAK1 antibody can be used for detection of DRAK1

> by Western blot at 1 μ g/mL. An approximately 50 kDa band can be detected. Antibody can also be used for immunocytochemistry starting at 2 μ g/mL. For immunofluorescence start at 10 µg/mL.Antibody

validated: Western Blot in human samples; Immunocytochemistry in human samples and Immunofluorescence in human samples. All other

applications and species not yet tested.

Isotype IgG

Clonality Polyclonal

MW 50 kDa


Uniprot ID Q9UEE5

NCBI

Dilution Range DRAK1 antibody can be used for detection of DRAK1

> by Western blot at 1 µg/mL. An approximately 50 kDa band can be detected. Antibody can also be used for immunocytochemistry starting at 2 ug/m


www.biorbyt.com



Western blot analysis of DRAK1 in (A) MO...



Immunocytochemistry staining of MOLT4 ce...



Immunofluorescence of DRAK1 in Molt cell...