



Product Datasheet

BMF Antibody (orb1240096)




www.biorbyt.com


A B

15-8-3-

> Western blot analysis of Bmf expression ...



Immunofluorescence of Bmf in human kidne...



Immunofluorescence of Bmf in Human Kidne...

Descriptionnts. BMF Antibody

Species/Host Rabbit

Reactivity Human

Conjugation Unconjugated

Tested ELISA, IF, IHC-P, WB

Applications

Immunogen Bmf antibody was raised with a synthetic peptide

corresponding to 14 amino acids near the carboxy terminus of human Bmf.The immunogen is located

within the last 50 amino acids of Bmf.

Target BMF

Preservatives Bmf Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/mL

Storage Bmf antibody can be stored at 4°C for three months

and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed

to prolonged high temperatures.

Note For research use only

Application notes Bmf antibody can be used for detection of Bmf by

Western blot at 2.5 and 5 µg/mL. A band at

approximately 25 kDa can be detected. Antibody can also be used for immunohistochemistry starting at 10

 μ g/mL. For immunofluorescence start at 10

μg/mL.Antibody validated: Western Blot in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype IgG

Clonality Polyclonal

MW 25 kDa

Uniprot ID Q96LC9

NCBI NP_277038

Dilution Range Bmf antibody can be used for detection of Bmf by

Western blot at 2.5 and 5 $\mu g/mL$. A band at

approximately 25 kDa can be detected. Antibody can also be used for immunohistochemistry starting at 10

ua/ml For immunofluorocconco ctart at 10

Carolina < br > 27709. United States

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North