

Product Datasheet

BIK Antibody (orb1240107)

Description

BIK Antibody

Species/Host

Rabbit

Reactivity

Human, Mouse

Conjugation

Unconjugated

Tested

ELISA, ICC, IF, WB

Applications

Immunogen

BIK antibody was raised against a 15 amino acid synthetic peptide from near the amino terminus of human BIK. The immunogen is located within the first 50 amino acids of Bik.

Target

BIK

Preservatives

Bik Antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/mL

Storage

Bik antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note

For research use only

Application notes

BIK antibody can be used for the detection of BIK by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunocytochemistry starting at 1 µg/mL. For immunofluorescence start at 10 µg/mL. Antibody validated: Western Blot in human samples; Immunocytochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype

IgG

Clonality

Polyclonal

MW

Predicted: 18 kDa Observed: 22 kDa

Uniprot ID


[Q13323](#)

NCBI


[CAG30276](#)

Dilution Range


BIK antibody can be used for the detection of BIK by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunocytochemistry starting at 1 µg/mL. For immunofluorescence start at 10 µg/mL. Antibody validated: Western Blot in human samples;



Western blot analysis of BIK in Jurkat c...



Immunocytochemistry of BIK in Jurkat cel...



Immunofluorescence of Bik in Jurkat cell...