

www.biorbyt.com


Product Datasheet

BCL2L10 Antibody (orb1240122)

biorbyt

Descriptionnts.	BCL2L10 Antibody
Species/Host	Rabbit
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	ELISA, ICC, WB
Immunogen	Bcl-B antibody was raised against a 16 amino acid synthetic peptide from near the amino terminus of human Bcl-B.The immunogen is located within amino acids 90 - 140 of Bcl-B.
Target	BCL2L10
Preservatives	Bcl-B Antibody is supplied in PBS containing 0.02% sodium azide.
Form/Appearance	Liquid
Concentration	1 mg/mL
Storage	Bcl-B antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Note	For research use only
Application notes	Bcl-B antibody can be used for the detection of Bcl- B by Western blot at $1 - 2 \mu g/mL$. Despite its predicted molecular weight, Bcl-B is often at higher molecular weights, presumably due to post- translational modifications. Antibody can also be used for immunocytochemistry starting at 10 $\mu g/mL$. Antibody validated: Western Blot in human samples and Immunocytochemistry in human samples. All other applications and species not yet tested.
lsotype	IgG
Clonality	Polyclonal
MW	Predicted: 22 kDa Observed: 26 kDa
Uniprot ID	Q9HD36
NCBI	NP_065129
Dilution Range	Bcl-B antibody can be used for the detection of Bcl- B by Western blot at $1 - 2 \mu g/mL$. Despite its

www.biorbyt.com



АВ

Western blot analysis of Bcl-B in Jurkat...



Immunocytochemistry of Bcl-B in Jurkat c...

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709. United States Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

samples. All other applications and species not yet tested

prodicted molecular weight Rel R is often at higher