

## Product Datasheet

### XIAP Antibody (orb1240298)

## Description

## XIAP Antibody

### Species/Host

Rabbit

### Reactivity

Human, Mouse

### Conjugation

Unconjugated

### Tested

ELISA, IF, IHC-P, WB

### Applications

### Immunogen

XIAP antibody was raised against a synthetic peptide corresponding to 13 amino acids at the C-terminus of human XIAP. The immunogen is located within amino acids 420 - 470 of XIAP.

### Target

XIAP

### Preservatives

XIAP Antibody is supplied in PBS containing 0.02% sodium azide.

### Form/Appearance

Liquid

### Concentration

1 mg/mL

### Storage

XIAP antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

### Note

For research use only

### Application notes

XIAP antibody can be used for the detection of XIAP by Western blot at 0.5 to 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 10 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human and mouse samples and Immunofluorescence in human and mouse samples. All other applications and species not yet tested.

### Isotype

IgG

### Clonality

Polyclonal

### MW

Predicted: 55 kDa Observed: 53 kDa

### Uniprot ID


[P98170](#)

### NCBI


[NP\\_001158](#)

### Dilution Range


XIAP antibody can be used for the detection of XIAP by Western blot at 0.5 to 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 10 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human and mouse samples



Western blot analysis of XIAP in human k...



Immunohistochemical staining of human ki...



Immunofluorescence of XIAP in Human Kidn...