

Product Datasheet

KRT10 Antibody (orb1252499)

Description

KRT10 Antibody

Species/Host

Mouse

Reactivity

Human, Mouse

Conjugation

Unconjugated

Tested

FC, IF, IHC-P, WB

Applications

Immunogen

Recombinant human KRT10 protein was used as the immunogen for the Keratin 10 antibody.

Target

KRT10

Preservatives

PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide

Form/Appearance

Liquid

Concentration

0.2 mg/mL

Storage

Aliquot and Store at 2-8°C. Avoid freeze-thaw cycles.

Note

For research use only

Application notes

Flow Cytometry: 0.5-1 ug/million cells in 0.1ml
Immunofluorescence: 0.5-1 ug/ml
Immunohistochemistry (FFPE): 0.1-0.2 ug/ml for 30 min at RT (1)Prediluted format : incubate for 30 min at RT (2)Optimal dilution of the Keratin 10 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Isotype

IgG1, kappa

Clonality


Monoclonal

Uniprot ID

P13645

Dilution Range


Flow Cytometry: 0.5-1 ug/million cells in 0.1ml
Immunofluorescence: 0.5-1 ug/ml
Immunohistochemistry (FFPE): 0.1-0.2 ug/ml for 30 min at RT (1)Prediluted format : incubate for 30 min at RT (2)Optimal dilution of the Keratin 10 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the



IHC testing of FFPE human skin with Kera...



IHC testing of FFPE human skin with Kera...



IHC testing of FFPE human bladder carcin...