


Product Datasheet

CXCL12 Antibody (orb1272312)

Description

CXCL12 Antibody

Species/Host	Goat
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	ELISA, WB
Immunogen	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hSDF-1- β (human Stromal Cell-Derived Factor-1-beta).
Target	CXCL12
Form/Appearance	Lyophilized
Concentration	batch dependent
Storage	SDF-1B antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.
Note	For research use only
Clonality	Polyclonal
Uniprot ID	P48061
NCBI	P48061
Dilution Range	ELISA:To detect hSDF-1- β by direct ELISA (using 100 μ L/well antibody solution) a concentration of at least 0.5 μ g/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hSDF-1- β . Sandwich:To detect hSDF-1- β by sandwich ELISA (using 100 μ L/well antibody solution) a concentration of 0.5 - 2.0 μ g/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human SDF-1- β as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hSDF-1- β . Western Blot:To detect hSDF-1- β by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hSDF-1- β is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Expiration Date	12 months from date of receipt.



To detect Human SDF-1-beta by sandwich E...



To detect Human SDF-1-beta by Western Bl...



To detect Human SDF-1-beta by Western Bl...