

Product Datasheet

CLEC11A Antibody (Biotin) (orb1272317)



www.biorbyt.com

Description of CLEC11A Antibody (Biotin)

Species/Host Goat

Reactivity Human

Conjugation Biotin

Tested ELISA, WB

Applications

Immunogen Produced from sera of goats pre-immunized with highly

pure (>98%) recombinant hSCGF-β (human SCGF-β).

Target CLEC11A

Form/Appearance Lyophilized

Concentration batch dependent

Storage SCGF-B antibody is stable for at least 2 years from date

of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C.

Avoid repeated freeze-thaw cycles.

Note For research use only

Application notes ELISA:Sandwich:To detect hSCGF-β by sandwich ELISA

(using 100 μ L/well antibody solution) a concentration of 0.25 - 1.0 μ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Human SCGF- β (XP-5268) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hSCGF- β . Western Blot:To detect hSCGF- β by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hSCGF- β is 1.5 - 3.0 ng/lane, under either reducing or non-reducing

conditions.

Clonality Polyclonal

Uniprot ID Q9Y240

NCBI Q9Y240

Dilution Range ELISA:Sandwich:To detect hSCGF-β by sandwich ELISA

(using 100 μ L/well antibody solution) a concentration of 0.25 - 1.0 μ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Human SCGF- β (XP-5268) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hSCGF- β . Western Blot:To detect hSCGF- β by Western Blot analysis this antibody

To detect hSCGF-beta by Western Blot ana...



To detect hSCGF-beta by Western Blot ana...