

Product Datasheet

CCL18 Antibody (orb1272359)

Description

CCL18 Antibody

Species/Host

Rabbit

Reactivity

Human

Conjugation

Unconjugated

Tested

ELISA, NeA, WB

Applications

Immunogen

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMIP-4 (human Macrophage Inflammatory Protein-4).

Target

CCL18

Form/Appearance

Lyophilized

Concentration

batch dependent

Storage

MIP-4 antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.

Note

For research use only

Clonality

Polyclonal

Uniprot ID


[P55774](#)

NCBI

[P55774](#)

Dilution Range


Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hMIPu4 (100.0 ng/mL), a concentration of 8.0 - 10.0 µg/mL of this antibody is required. ELISA: To detect hMIP-4 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hMIP-4. Sandwich: To detect hMIP-4 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human MIP-4 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIP-4. Western Blot: To detect hMIP-4 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIP-4 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.



To detect hMIP-4 by Western Blot analysis...



To detect hMIP-4 by Western Blot analysis...



This antibody stained formalin-fixed, pa...