Product Datasheet

CCL4 Antibody (Biotin) (orb1272370)

|	
Description	CCL4 Antibody (Biotin)
Species/Host	Rabbit
Reactivity	Human
Conjugation	Biotin
Tested Applications	ELISA, WB
Immunogen	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hMIP-1-β (human MIP-1-β).
Target	CCL4
Form/Appearance	Lyophilized
Concentration	batch dependent
Storage	MIP-1B antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.
Note	For research use only
Application notes	ELISA:Sandwich:To detect hMIP-1β by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Human MIP-1β (XP-5235) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIP-1β.Western Blot:To detect hMIP-1-β by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIP-1-β is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Clonality	Polyclonal
Uniprot ID	P13236
NCBI	P13236
Dilution Range	ELISA:Sandwich:To detect hMIP-1β by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Human MIP-1β (XP-5235) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIP-1β.Western Blot:To detect hMIP-1-β by Western Blot analysis this antibody can be

To detect Human MIP-1-beta by sandwich E...

To detect Human MIP-1-beta by Western Bl...

To detect Human MIP-1-beta by Western Bl...

Expiration Date	12 months from date of receipt.