

## Product Datasheet

### IL9 Antibody (orb1272436)

## Description

## IL9 Antibody

### Species/Host

Rabbit

### Reactivity

Human

### Conjugation

Unconjugated

### Tested

ELISA, NeA, WB

### Applications

### Immunogen

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-9 (human Interleukin-9).

### Target

IL9

### Form/Appearance

Lyophilized

### Concentration

batch dependent

### Storage

IL-9 antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.

### Note

For research use only

### Clonality

Polyclonal

### Uniprot ID

**P15248**

### NCBI


**P15248**

### Dilution Range


Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hIL-9 (0.6 ng/mL), a concentration of 0.15 - 0.025 µg/mL of this antibody is required. ELISA: To detect hIL-9 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hIL-9. Sandwich: To detect hIL-9 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human IL-9 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-9. Western Blot: To detect hIL-9 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-9 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

### Expiration Date


12 months from date of receipt.



To detect Human IL-9 by sandwich ELISA (...)



To detect Human IL-9 by Western Blot ana...



To detect Human IL-9 by Western Blot ana...