

www.biorbyt.com

Product Datasheet

CXCL3 Antibody (orb1272515)

biorbyt

Description ^{nts.}	CXCL3 Antibody	
Species/Host	Rabbit	1008-0
Reactivity	Human	00 100 100 100 100
Conjugation	Unconjugated	
Tested Applications	ELISA, NeA, WB	
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hGRO-gamma (human GRO- gamma).	
Target	CXCL3	-
Form/Appearance	Lyophilized	
Concentration	batch dependent	
Storage	Gro-gamma antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
Note	For research use only	۱
Clonality	Polyclonal	
Uniprot ID	P19876	
NCBI	P19876	
Dilution Range	Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hGRO-gamma (100 ng/mL), a concentration of $1.0 - 2.0 \mu$ g/mL of this antibody is required. ELISA:To detect hGRO-gamma by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of $0.2 - 0.4$ ng/well of recombinant hGRO-gamma.Sandwich: To detect hGRO- γ by sandwich ELISA (using 100 µL/well antibody solution) a concentration of $0.5 - 2.0 \mu$ g/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human GRO- γ as a detection antibody, allows the detection of at least $0.2 -$ 0.4 ng/well of recombinant hGRO- γ . Western Blot:To detect hGRO-gamma by Western Blot analysis this antibody can be used at a concentration of $0.1 - 0.2 \mu$ g/mL. Used in	Ĭ

www.biorbyt.com Ráitit kei Annar 190-yanna



To detect Human GROgamma by sandwich EL...



To detect Human GROgamma by Western Blo ...



To detect Human GROgamma by Western Blo ...

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280

conjunction with compatible secondary reagents the detection limit for recombinant hGRO-gamma is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

> 68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709. United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558