





## Product Datasheet

### FGF10 Antibody (orb1272550)

Description	FGF10 Antibody	
Species/Host	Goat	
Reactivity	Human	
Conjugation	Unconjugated	
Tested Applications	ELISA, WB	
Immunogen	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hFGF-10 (human Fibroblast Growth Factor-10).	This antibody stained formalin-fixed par...
Target	FGF10	
Form/Appearance	Lyophilized	
Concentration	batch dependent	
Storage	FGF-10 antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
Note	For research use only	
Clonality	Polyclonal	
Uniprot ID	<a href="#">O15520</a>	
NCBI	<a href="#">O15520</a>	
Dilution Range	ELISA: To detect hFGF-10 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hFGF-10. Sandwich: To detect hFGF-10 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human FGF-10 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hFGF-10. Western Blot: To detect hFGF-10 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hFGF-10 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.	This antibody stained formalin-fixed par...
Expiration Date	12 months from date of receipt.	