

Product Datasheet

CCL26 Antibody (orb1272558)

Description

CCL26 Antibody

Species/Host

Goat

Reactivity

Human

Conjugation

Unconjugated

Tested

ELISA, NeA, WB

Applications

Immunogen

Produced from sera of goats pre-immunized with highly pure (>98%) recombinant human Eotaxin-3.

Target

CCL26

Form/Appearance

Lyophilized

Concentration

batch dependent

Storage

Eotaxin-3 antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.

Note

For research use only

Clonality

Polyclonal

Uniprot ID


[Q9Y258](#)

NCBI


[Q9Y258](#)

Dilution Range


Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of human eotaxin-3 (100.00 ng/mL), a concentration of 5.0 - 7.0 µg/mL of this antibody is required. ELISA: To detect human eotaxin-3 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant human eotaxin-3. Sandwich: To detect hEotaxin-3 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human Eotaxin-3 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hEotaxin-3. Western Blot: To detect human eotaxin-3 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant human eotaxin-3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.



To detect hEotaxin-3 by sandwich ELISA (...)



To detect hEotaxin-3 by Western Blot ana...



To detect hEotaxin-3 by Western Blot ana...