


## Product Datasheet

### TNFSF13B Antibody (orb1272595)


## Description

## TNFSF13B Antibody

<b>Species/Host</b>	Goat
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Tested Applications</b>	ELISA, WB
<b>Immunogen</b>	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hBAFF (human BAFF).
<b>Target</b>	TNFSF13B
<b>Form/Appearance</b>	Lyophilized
<b>Concentration</b>	batch dependent
<b>Storage</b>	BAFF antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.
<b>Note</b>	For research use only
<b>Application notes</b>	ELISA:To detect hBAFF by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hBAFF. Western Blot:To detect hBAFF by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hBAFF is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
<b>Clonality</b>	Polyclonal
<b>Uniprot ID</b>	<a href="#">Q9Y275</a>
<b>NCBI</b>	<a href="#">Q9Y275</a>
<b>Dilution Range</b>	ELISA:To detect hBAFF by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hBAFF. Western Blot:To detect hBAFF by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hBAFF is 1.5 - 3.0 ng/lane, under either



To detect hBAFF by sandwich ELISA (using...



To detect hBAFF by Western Blot analysis...