

## **Product Datasheet**

IL19 Antibody (Biotin) (orb1272763)



## www.biorbyt.com

**Description**nts. IL19 Antibody (Biotin)

Species/Host Rabbit

**Reactivity** Human

**Conjugation** Biotin

Tested ELISA, WB

**Applications** 

**Immunogen** Produced from sera of rabbits pre-immunized with

highly pure (>98%) recombinant hIL-19. Human IL-19

specific antibody was purified by affinity

chromatography and then biotinylated.ced from sera of

rabbits pre-immunized with highly pure (>98%) recombinant hIL-19. Human IL-19 specific antibody was

purified by affinity chromatography employing immobilized hlL-19 matrix.

Target IL19

Form/Appearance Lyophilized

**Concentration** batch dependent

**Storage** IL-19 antibody is stable for at least 2 years from date of

receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid

repeated freeze-thaw cycles.

**Note** For research use only

**Application notes** ELISA:Sandwich:To detect hIL-19 by sandwich ELISA

(using 100  $\mu$ L/well antibody solution) a concentration of 0.25 - 1.0  $\mu$ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our polyclonal Anti-Human IL-19 as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-19. Western Blot:To detect hIL-19 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2  $\mu$ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-19 is 1.5 - 3.0 ng/lane, under either

reducing or non-reducing conditions.

**Clonality** Polyclonal


Uniprot ID Q9UHD0

NCBI Q9UHD0

**Dilution Range** ELISA:Sandwich:To detect hIL-19 by sandwich ELISA

(using 100  $\mu\text{L/well}$  antibody solution) a concentration of

0.25 - 1.0 μg/mL of this antibody is required. This



To detect hIL-19 by sandwich ELISA (usin...



To detect hIL-19 by Western Blot analysi...



To detect hIL-19 by Western Blot analysi...