

## Product Datasheet

### BAFF Antibody [Buffy-2] (orb1273961)

**Description**

BAFF Antibody [Buffy-2]

**Species/Host**

Rat

**Reactivity**

Human, Mouse

**Conjugation**

Unconjugated

**Tested**

FC, IHC, WB

**Applications**
**Immunogen**

BAFF (monoclonal Buffy-2) antibody was raised against recombinant human soluble BAFF (aa. 83-285).

**Preservatives**

PBS containing 0.02% sodium azide

**Form/Appearance**

Liquid

**Concentration**

batch dependent

**Storage**

BAFF antibody can be stored at 4°C, stable for one year.

**Note**

For research use only

**Application notes**

BAFF antibody can be used to recognize membrane-bound and soluble human and mouse (weak) BAFF by Flow Cytometry, Immunohistochemistry (frozen sections, paraffin sections) and Western Blot. Method: HEK 293T cells (5 x 10<sup>5</sup>) were mock transfected (thin line) or transfected with an expression plasmid enabling surface expression of full length hBAFF (thick line). Cells were incubated on ice for 30min in 50 µl FACS buffer (PBS, 5% fetal calf serum, 0.02% azide) containing 1 µg/mL of MAb to BAFF (Buffy-2) (FITC). After washing, cells were analyzed by flow cytometry. Method: Tissues are in paraffin sections. Sections are de-waxed and rehydrated, then cooked. Sections are stained in wet boxes, rehydrated and finally prepared as coverslip slides using Eukitt solution. Please inquire for complete protocol..

**Isotype**


IgM

**Clonality**

Monoclonal

**Uniprot ID**
**Q9Y275**
**Dilution Range**

BAFF antibody can be used to recognize membrane-bound and soluble human and mouse (weak) BAFF by Flow Cytometry, Immunohistochemistry (frozen sections, paraffin sections) and Western Blot. Method: HEK 293T cells (5 x 10<sup>5</sup>) were mock transfected (thin line) or transfected with an expression plasmid enabling surface expression of full length hBAFF (thick line). Cells were incubated on ice for 30min in 50 µl FACS buffer (PBS, 5% fetal calf serum, 0.02% azide) containing 1 µg/mL of MAb to BAFF (Buffy-2) (FITC). After washing, cells were



The immunogen is recombinant human solub...

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive&lt;br&gt;Research Triangle Park&lt;br&gt;Durham, North Carolina&lt;br&gt;27709. United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

**Expiration Date**

12 months from date of receipt.