

Product Datasheet

CD210 antibody (orb1293374)

Description

Rat monoclonal antibody to CD210

Species/Host

Rat

Reactivity

Mouse

Conjugation

Unconjugated

Tested

FC, IF

Applications
Immunogen

Soluble mIL-10R-FLAG-His

Target

IL10RA

Preservatives

Rat IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

Form/Appearance

Rat IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

Storage

Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Note

For research use only

Application notes

1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube. 2. Add appropriate amount of antibody to the bottom of flow tube mixing with the whole blood; incubate for 30 minutes at room temperature. 3. Add 2 ml RBC lysis buffer; incubate for 10 minutes after mixing; dissolve red blood cells. 4. Sample tube is set to 1000 rpm centrifugation for 5 minutes; discard the supernatant. 5. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 6. Add appropriate amount of fluorescent-labeled anti-rat IgGs and incubate for 20 minutes away from light at room temperature. 7. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 8. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Clonality

Monoclonal

Source

Rat

Uniprot ID
Q61727