

Product Datasheet

TCR beta antibody (FITC) (orb1293840)

Description

Hamster monoclonal antibody against TCR

Species/Host	Hamster
Reactivity	Mouse
Conjugation	FITC
Tested Applications	FC, IF
Immunogen	Affinity purified TCR from mouse DO-11.10 cells
Target	TCRB
Preservatives	Armenian Hamster IgG. Liquid in PBS, pH 7.3, and 0.02% sodium azide.
Form/Appearance	Armenian Hamster IgG. Liquid in PBS, pH 7.3, and 0.02% sodium azide.
Storage	Shipped and store at 4°C for one year. Do not freeze.
Note	For research use only
Application notes	1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube. 2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light. 3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells. 4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant. 5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant. 6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).
Clonality	Monoclonal
Source	Armenian Hamster
Entrez	21577
Dilution Range	10 µl / assay