

Product Datasheet

Myo1H antibody (orb131704)



www.biorbyt.com

250kDa 150kDa

50kDa 37kDa

> 25kDa 20kDa 15kDa

> 10kDa

Western blot analysis of NIH3T3 lysate (...



Immunohistochemical staining of Human Ki...

Descriptionnts. Goat polyclonal to Myo1H

Species/Host Goat

Reactivity Bovine, Human, Mouse, Porcine, Rat

Conjugation Unconjugated

Tested ELISA, IHC, WB

Applications

Target Myosin IH

Preservatives Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium

azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and

thawing.

Storage Aliquot and store at -20°C. Minimize freezing and

thawing.

Note For research use only

Application notes Approx 110kDa band observed in lysates of cell line

NIH3T3 (calculated MW of 118kDa according to NP_001158045.1). Recommended concentration: 0.1-0.3 µg/ml. An additional band of unknown identity was also consistently observed at 28kDa. This band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field-have any such results been reported with other antibodies/lysates Have any further splice variants/modified forms been reportedWB: Approx 110kDa band observed in lysates of cell line NIH3T3 (calculated MW of 118kDa according to NP 001158045.1). Recommended concentration: 0.1-0.3 µg/ml. An additional band of unknown identity was also consistently observed at 28kDa. This band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field have any such results been reported with other

antibodies/lysates Have any further splice variants/modified forms been reported

Protein Sequence EDQVYKDKNGQLR

Clonality Polyclonal

MW 118

Entrez 283446

Dilution Range ELISA: 1:128000, WB: 0.1-0.3 μg/ml, IHC-P: 5 μg/ml

Expiration Date 12 months from date of receipt.

Carolina

27709. United States