




Product Datasheet

HSD17B1 Antibody (orb137855)




www.biorbyt.com

Immunohistochemical analysis of formalin...



Immunohistochemical analysis of formalin...



Confocal immunofluoroscence analysis of ...

Descriptionnts. HSD17B1 Antibody

Species/Host Rabbit

Reactivity Human, Rat

Conjugation Unconjugated

Tested ICC. IHC. WB

Applications

A synthetic peptide corresponding to a sequence at **Immunogen**

> the N-terminus of human HSD17B1 (29-43aa QSFKVYATLRDLKTQ), different from the related rat

and mouse sequences by one amino acid.

Form/Appearance Lyophilized

Concentration Adding 0.2 ml of distilled water will yield a

concentration of 500 µg/ml.

Storage Store at -20°C for one year from date of receipt.

> After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Note For research use only

Application notes Tested Species: In-house tested species with positive

> results. Predicted Species: Species predicted to be fit for the product based on sequence similarities. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users. . Add 0.2ml of distilled

water will yield a concentration of 500ug/ml.

Isotype Rabbit IgG

Clonality Polyclonal

мw 34950 MW

P14061 **Uniprot ID**

Dilution Range Western blot, 0.1-0.5µg/ml, Human, Rat

Immunohistochemistry (Paraffin-embedded Section),

0.5-1µg/ml, Human, Rat, By Heat

Immunocytochemistry , 0.5-1µg/ml, Human

Expiration Date 12 months from date of receipt.

Carolina < br > 27709. United States

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North