

## **Product Datasheet**

NAK/TBK1 Antibody (orb137920)



Descriptionnts.

## www.biorbyt.com

NAK/TBK1 Antibody

Species/Host Rabbit

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested IHC, WB

**Applications** 

**Immunogen** A synthetic peptide corresponding to a sequence at

the C-terminus of human NAK (577-590aa

YNEEQIHKFDKQKL), identical to the related rat and

mouse sequences.

Form/Appearance Lyophilized

**Concentration** Adding 0.2 ml of distilled water will yield a

concentration of 500 µg/ml.

**Storage** Store at -20°C for one year from date of receipt.

After reconstitution, at  $4^{\circ}\text{C}$  for one month. It can also be aliquotted and stored frozen at  $-20^{\circ}\text{C}$  for six

months. Avoid repeated freeze-thaw cycles.

**Note** For research use only

**Application notes** WB: The detection limit for TBK1 is approximately

3ng/lane under reducing conditions. Tested Species:

In-house tested species with positive results.

Predicted Species: Species predicted to be fit for the product based on sequence similarities. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users. . Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Isotype Rabbit IgG

**Clonality** Polyclonal

**MW** 83642 MW


Uniprot ID Q9UHD2

**Dilution Range** Immunohistochemistry (Paraffin-embedded Section),


0.5-1µg/ml, Human, Rat, Mouse, By HeatWestern

blot, 0.1-0.5µg/ml, Human, Rat, Mouse

**Expiration Date** 12 months from date of receipt.



Immunohistochemical analysis of formalin...



Western blot analysis of Lane 1: Rat Tes...



Immunohistochemical analysis of formalin...

68 TW Alexander Drive<br/>br>Research Triangle Park<br/>br>Durham, North