

Product Datasheet

Rat IgG1 kappa Isotype Control-APC labled Antibody (orb1881028)

Description

Rat IgG1 kappa Isotype Control-APC

control in flow cytometry and ot...

Conjugation

APC

Tested

FC, IF

Applications
Form/Appearance

Rat IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

Storage

Shipped and store at 4°C for one year. Do not freeze.

Note

For research use only

Application notes

1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube.
2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light.
3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells.
4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.
5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.
6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Source

Rat

Dilution Range

10 µl / assay

Expiration Date

12 months from date of receipt.