

Product Datasheet

SKAP2 antibody (orb18961)

Description

Goat polyclonal antibody to SKAP2

Species/Host

Goat

Reactivity

Bovine, Canine, Human, Mouse, Rat

Conjugation

Unconjugated

Tested

ELISA, WB

Applications
Target

SCAP2 / PRAP

Preservatives

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Note

For research use only

Application notes

ELISA: Peptide ELISA: antibody detection limit dilution 1:32000. WB: Approx 38kDa band observed in lysates of cell line Jurkat (calculated MW of 41.2kDa according to NP_003921.2). In transfected HEK293 transiently expressing SKAP2 a band of approx. 45kDa is observed. This band is not observed in the non-transfected HEK293. The calculated Recommended concentration: 0.5-2 µg/ml. An additional band of unknown identity was also consistently observed in Jurkat at 28kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates Have any further splice variants/modified forms been reported

Protein Sequence

GLVPKAYIMEMYDI

Clonality

Polyclonal

MW

41.2

Entrez


8935

Dilution Range


ELISA: 1:32000, WB: 0.5-2 µg/ml

Expiration Date

12 months from date of receipt.

Western blot analysis of Jurkat lysate u...



Western blot analysis of HEK293 lysate u...