

Product Datasheet

CD2, CD20 antibody (FITC, PE) (orb198142)

Description

Mouse Polyclonal antibody to CD2, CD20

Species/Host

Mouse

Reactivity

Human

Conjugation

FITC/RPE

Tested

FACS

Applications
Immunogen

CD2=Derived from the hybridization of mouse Sp2/0 myeloma cells with spleen cells from BALB/c mice immunized with t lymphocytes activated by mixed lymphocyte culture.CD20=Derived from the hybridization of mouse Sp2/0 myeloma cells with spleen cells from BALB/c mice immunized with the LB lymphoblastoid cell line.

Preservatives

Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein

Form/Appearance

Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein

Concentration

Titered for flow cytometry

Storage

Product should be stored at 4-8°C. DO NOT FREEZE

Note

For research use only

Application notes

PBMC: Add10 µl of MAB/10⁶ PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2° to 8° C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature 20° C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturer's instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.

Isotype

IgG2a (F)/IgG1 (PE)

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709. United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

Expiration Date

12 months from date of receipt.