

## **Product Datasheet**

APE1/APEX1 Antibody (orb215883)



## www.biorbyt.com

	· ·	Www.h
Description <sup>nts.</sup>	APE1/APEX1 Antibody	1 3.4K 3.9 多枝
Species/Host	Rabbit	
Reactivity	Human, Mouse, Rat	8.
Conjugation	Unconjugated	0110 110 110 110
Tested Applications	FC, ICC, IF, IHC, WB	Flow Cytometry
lmmunogen	E.coli-derived human APE1 recombinant protein (Position: P2-L318). Human APE1 shares 94% and 93% amino acid (aa) sequences identity with mouse and rat APE1, respectively.	analysis of U937 cells us
Form/Appearance	Lyophilized	100KD —
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 $\mu\text{g/ml}$ .	70KD — 55KD —
Storage	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.	35KD — 25KD — 15KD — WB analysis of
Note	For research use only	APEX1 using anti-APEX1
Application notes	WB: The detection limit for APE1 is approximately 0.25ng/lane under reducing conditions. Tested Species: In-house tested species with positive results. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users Add 0.2ml of distilled water will yield a concentration of 500ug/ml.	an  1 2 3 4 5 6 7 8 9  116KD -  97KD -  58KD -  40KD -  29KD -  20KD -
Isotype	Rabbit IgG	14KD <b>-</b>
Clonality	Polyclonal	WB analysis of APEX1 using
MW	35555 MW	anti-APEX1 an
Uniprot ID	P27695	
Dilution Range	Western blot, 0.1-0.5μg/ml, Human, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 0.5-1μg/ml, Human, Mouse, Rat, By Heat Immunocytochemistry/Immunofluorescence, 2 μg/ml, Human Immunofluorescence, 2μg/ml, Human Flow Cytometry, 1-3μg/1x106 cells, Human	

**Expiration Date** 

12 months from date of receipt.