



Product Datasheet

FHIT Antibody (orb215935)



www.biorbyt.com

Description ^{nts.}	FHIT Antibody
-----------------------------	---------------

Species/Host Rabbit

Reactivity Human, Rat

Conjugation Unconjugated

Tested FC, ICC, IF, IHC, WB

Applications

Immunogen E.coli-derived human FHIT recombinant protein (Position: M1-

Q147). Human FHIT shares 90% and 87% amino acid (aa) sequences identity with mouse and rat FHIT, respectively.

Form/Appearance Lyophilized

Concentration Adding 0.2 ml of distilled water will yield a concentration of 500

μg/ml.

Storage Store at -20°C for one year from date of receipt. After

reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated

freeze-thaw cycles.

Note For research use only

Application notes WB: The detection limit for FHIT is approximately 0.25ng/lane

under reducing conditions. Tested Species: In-house tested species with positive results. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users. . Add 0.2ml of distilled water will yield a

concentration of 500ug/ml.

Isotype Rabbit IgG

Clonality Polyclonal

MW 16858 MW


Uniprot ID P49789

Dilution Range Western blot, 0.1-0.5μg/ml, Human Immunohistochemistry


(Paraffin-embedded Section), 0.5-1µg/ml, Human, By Heat Immunocytochemistry/Immunofluorescence, 2µg/ml, Human

Flow Cytometry, 1-3µg/1x106 cells, Human


Expiration Date 12 months from date of receipt.



Flow Cytometry analysis of U20S cells us...



WB analysis of FHIT using anti-FHIT anti...



IF analysis of FHIT using anti-FHIT anti...

Carolina

27709. United States

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North