

Product Datasheet

TG Antibody Cocktail / Thyroglobulin (orb248559)

Description

Thyroglobulin is a 660kDa dimeric pre-protein



~300kDa ...

Species/Host

Mouse

Reactivity


Human, Mouse, Rat

Conjugation

Unconjugated

Tested

IHC-P

Applications


Immunohistochemical
analysis of formalin...

Immunogen

Human thyroid follicular cells were used as the immunogen for this Thyroglobulin antibody.

Preservatives

0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin (US sourced) and 0.05% sodium azide

Storage

Store the Thyroglobulin antibody cocktail at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

Note

For research use only

Application notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titrated up or down for optimal performance.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Formula

0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide

Isotype

Mouse IgG1, kappa

Clonality

Monoclonal

Purity

Protein G affinity chromatography

Hazard
Information

This Thyroglobulin antibody is available for research use only.

Entrez

7038

Dilution Range

Immunohistochemistry (FFPE): 0.1-0.2ug/ml for 30 min at RT