


Product Datasheet

Human Ig light chain Lambda antibody (orb248958)

Description

Mouse monoclonal antibody to Human Ig light chain



Flow
cytometric
analysis of
human
periph...

Species/Host

Mouse

Reactivity

Human

Conjugation

Unconjugated

Tested

FC, IF

Applications

Immunogen

Native purified human Ig light chain Lambda.

Preservatives

Mouse IgG1. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

Form/Appearance

Mouse IgG1. Liquid in PBS, pH 7.3, and 0.02% sodium azide.

Storage

Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Note

For research use only

Application notes

1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube. 2. Add appropriate amount of antibody to the bottom of flow tube mixing with the whole blood; incubate for 30 minutes at room temperature. 3. Add 2 ml RBC lysis buffer; incubate for 10 minutes after mixing; dissolve red blood cells. 4. Sample tube is set to 1000 rpm centrifugation for 5 minutes; discard the supernatant. 5. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 6. Add appropriate amount of fluorescent-labeled anti-mouse IgGs and incubate for 20 minutes away from light at room temperature. 7. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 8. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Clonality

Monoclonal

Source

Mouse

Expiration Date

12 months from date of receipt.