

Product Datasheet

PDPK1 Antibody (orb316598)



iorbyt.com

| liorbyt | | www.bior |
|------------------------|---|---|
| Descriptionnts. | PDPK1 Antibody | 130KD – 1 2 3 4 5 6 |
| Species/Host | Rabbit | 70KD |
| Reactivity | Human, Mouse, Rat | 55KD 35KD |
| Conjugation | Unconjugated | 25KD – 15KD – |
| Tested Applications | IHC, WB | WB analysis of PDPK1 using |
| Immunogen | A synthetic peptide corresponding to a sequence at the C-terminus of human PDPK1 (524-556aa | anti-PDPK1 an |
| | YLMDPSGNAHKWCRKIQEVWRQRYQSHPDAAVQ), different fro the related mouse and rat sequences by two amino acids. | |
| Form/Appearance | Lyophilized | |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 $\mu\text{g/ml}.$ | |
| Storage | Store at -20°C for one year from date of receipt. After | |
| | reconstitution, at 4°C for one month. It can also be aliquotte and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. | d IHC analysis of PDPK1 using anti-PDPK1 a |
| Note | For research use only | |
| Application notes | Tested Species: In-house tested species with positive results By Heat: Boiling the paraffin sections in 10mM citrate buffer pH6.0, for 20mins is required for the staining of | |



IHC analysis of PDPK1 using anti-PDPK1 a...

| Isotype Rak | bit IgG |
|--------------------|---------|
|--------------------|---------|

Clonality Polyclonal

MW 63152 MW

Uniprot ID 015530

Dilution Range Immunohistochemistry (Paraffin-embedded Section), 0.5-

1µg/ml, By HeatImmunohistochemistry (Frozen Section), 0.5-

formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users. . Add 0.2ml of distilled water will yield a concentration of

 $1\mu g/ml$ Western blot, $0.1-0.5\mu g/ml$

Expiration Date 12 months from date of receipt.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

500ug/ml.