

## Product Datasheet

### Acetyl Cholinesterase antibody (orb344207)

**Description**

Acetyl Cholinesterase antibody

**Species/Host**

Goat

**Reactivity**

Bovine

**Conjugation**

Unconjugated

**Tested**

ELISA, IP, WB

**Applications**
**Immunogen**

Acetyl Cholinesterase [Bovine Erythrocytes]

**Preservatives**

0.01% (w/v) Sodium Azide

**Form/Appearance**

Liquid (sterile filtered)

**Concentration**

1.0 mg/ml

**Storage**

Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

**Note**

For research use only

**Application notes**

Anti-Acetyl Cholinesterase Antibody has been tested by western blot and is suitable to be assayed against 1.0 µg of Acetyl Cholinesterase [Bovine Erythrocytes] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) code #605-4302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,200 to 1:5,000 of the reconstitution concentration is suggested for this product.

**Isotype**

IgG

**Clonality**

Polyclonal

**Purity**

Anti-Acetyl Cholinesterase is an IgG fraction antibody purified from

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive&lt;br&gt;Research Triangle Park&lt;br&gt;Durham, North Carolina&lt;br&gt;27709, United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat