

Product Datasheet

Swine IgG (H&L) antibody (TRITC) (orb347970)



www.biorbyt.com

Descriptionnts. Swine IgG (H&L) antibody (TRITC)

Species/Host Rabbit

Porcine Reactivity

TRITC Conjugation

Tested

DOT, FC, FLISA, IF, WB

Applications

Swine IgG whole molecule **Immunogen**

0.01% (w/v) Sodium Azide **Preservatives**

Form/Appearance Lyophilized

Concentration 2.0 mg/mL

Store vial at 4° C prior to restoration. For Storage

> extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Note For research use only

Application notes Anti-Swine IgG (H&L) Rhodamine Antibody

has been tested by dot blot and western

blot and is designed for

immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Isotype IgG

Clonality Polyclonal

Purity This product was prepared from

monospecific antiserum by

immunoaffinity chromatography using Swine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Swine IgG and Swine Serum.

Dilution Range FLISA: 1:10,000 - 1:50,000, FC: 1:500 -

1:2,500, IF: 1:1,000 - 1:5,000