

Product Datasheet

F(ab')2 Bovine IgG (H&L) antibody (orb348074)



www.biorbyt.com

Descriptionnts. F(ab')2 Bovine IgG (H&L) antibody

Species/Host Rabbit

Reactivity Bovine

Conjugation Unconjugated

Tested ELISA, IHC, WB

Applications

Immunogen Bovine IgG whole molecule

Preservatives 0.01% (w/v) Sodium Azide

Form/Appearance Liquid (sterile filtered)

Concentration 1.0 mg/mL

Storage Store vial at 4° C prior to opening. This

product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and

thawing.

Note For research use only

Application notes Suitable for immunomicroscopy and flow

cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most

applications.

Isotype IgG F(ab')2

Clonality Polyclonal

Purity This product was prepared from

monospecific antiserum by

immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
br>27709. United States

 $\begin{tabular}{ll} Email: info@biorbyt.com & Phone: +1 (415) 906-5211 & Fax: +1 (415) 651-8558 \\ \end{tabular}$

Dilution Range ELISA: 1:20,000-1:300,000, IHC: 1:1,000-