

Product Datasheet

**F(ab')₂ Goat IgG (H&L) antibody (Peroxidase)
(orb348117)**

Description

F(ab')₂ Goat IgG (H&L) antibody

Species/Host

Rabbit

Reactivity

Goat

Conjugation

HRP

Tested

ELISA, IHC, WB

Applications
Immunogen

Anti-Goat IgG whole molecule was produced by repeated immunization with Goat IgG whole molecule in rabbit.

Preservatives

0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Form/Appearance

Lyophilized

Concentration

1.0 mg/mL

Storage

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Note

For research use only

Application notes

F(ab')₂ Anti-Goat IgG Peroxidase Antibody has been tested by ELISA and is assayed against 1.0 ug of Goat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:50,000 is suggested for this product. This secondary conjugated antibody anti-goat is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.

Isotype

IgG F(ab')₂
Clonality

Polyclonal

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709, United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

to remove any unwanted reactivities,
pepsin digestion and chromatographic