

## Product Datasheet

**F(ab')<sub>2</sub> Rabbit IgG F(c) antibody (Biotin)  
(orb348313)**

**Description**

F(ab')<sub>2</sub> Rabbit IgG F(c) antibody (Biotin)

**Species/Host**

Goat

**Reactivity**

Rabbit

**Conjugation**

Biotin

**Tested**

ELISA, IHC, WB

**Applications**
**Immunogen**

Rabbit IgG F(c) fragment

**Preservatives**

0.01% (w/v) Sodium Azide

**Form/Appearance**

Lyophilized

**Concentration**

1.0 mg/mL

**Storage**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Note**

For research use only

**Application notes**

F(ab')<sub>2</sub> Anti-Rabbit IgG F(c) Biotin Antibody has been tested by SDS-PAGE and western blot and is assayed against 1.0 ug of Rabbit IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:12,000 to 1:60,000 of the reconstitution concentration is suggested for this product. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.

**Isotype**

IgG F(ab')<sub>2</sub>
**Clonality**

Polyclonal

**Purity**

F(ab') fragment biotin conjugated secondary antibody was prepared from

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive&lt;br&gt;Research Triangle Park&lt;br&gt;Durham, North Carolina&lt;br&gt;27709, United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

pepsin digestion and chromatographic separation. Assay by