

Product Datasheet

A431 Whole Cell Lysate (orb348665)



www.biorbyt.com

Descriptionnts. A431 Whole Cell Lysate

Conjugation Unconjugated

Tested
Applications

SDS-PAGE, WB

Preservatives Preservative: None. Stabilizer: 10% (v/v)

Glycerol. 1X SDS-PAGE Sample Buffer (62.5 mM Tris HCl, 2% SDS, 10% Glycerol and 0.005% bromophenol blue, pH 6.8)

Form/Appearance Liquid

Concentration 1.0 mg/ml

Storage Store A431 Whole Cell Lysate at -70° C or

COLDER. For extended storage, whole cell lysate to minimize freeze/thaw cycles.

Note For research use only

Application notes ready-to-use A431 Whole Cell Lysate are

especially prepared as positive controls for separation by SDS-PAGE and subsequent western blot analysis. Lysates are prepared in denaturing buffer WITHOUT dissociating agents (i.e. no 2-mercaptoethanol or dithiothreitol has been added). Heat lysate to 95°C for 5 minutes and rapidly cool. If dissociating conditions are desired, add reducing agent prior to heating. The recommended loading volume per lane is 10-

recommended loading volume per lane is 10-20 µl depending on the size format of your

gel.

Purity A423 cells were grown in Dulbecco's

medium supplemented with 10% fetal bovine serum. Cells were washed with PBS and then incubated on ice in modified RIPA buffer, containing 150 mM sodium chloride, 50 mM Tris HCl, pH 7.4, 1 mM EDTA, 1.0% NP-40, 0.5% sodium deoxycholic acid, 0.1% SDS and 0.01% (w/v) sodium azide to lyse the cells. Protein integrity was ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and serine proteases as well as aminopeptidases (0.1 mM AEBSF HCl, 0.08 μM Aprotinin, 5 μM Bestatin, 1.5 μM E-64, 2 μM Leupeptin Hemisulfate, 1 μM Pepstatin A). Phosphatase inhibitors 1 mM NaF and 1 mM Na3VO4 were also added. Cell debris was removed by centrifugation. Protein