

Product Datasheet

HeLa Whole Cell Lysate TNFa Stimulated (orb348673)



www.biorbyt.com

Description nts. HeLa Whole Cell Lysate TNFa

Conjugation Unconjugated

Tested Applications

SDS-PAGE, WB

Preservatives Preservative: None. Stabilizer: 10%

(v/v) Glycerol. 1X SDS-PAGE Sample Buffer (62.5 mM Tris HCl, 2% SDS, 10% Glycerol and 0.005% bromophenol

blue, pH 6.8)

Form/Appearance Liquid

Concentration 0.97 mg/ml

Storage Store vial at -70° C or COLDER. For

extended storage, aliquot contents to

minimize freeze/thaw cycles.

Note For research use only

Application notes ready-to-use lysates are especially

prepared as positive controls for separation by SDS-PAGE and subsequent western blot analysis. Lysates are prepared in denaturing buffer WITHOUT dissociating agents

(i.e. no 2-mercaptoethanol or dithiothreitol has been added). Heat lysate to 95°C for 5 minutes and rapidly cool. If dissociating conditions are desired, add reducing agent prior to heating. The recommended loading volume per lane is 10-20 µl depending

on the size format of your gel.

Purity The cells were grown in Dulbecco's

medium supplemented with 10% fetal bovine serum. Cells were treated with 0.2 μ g/ml TNF- α for 30 min. Cells were washed with PBS and then incubated

on ice in modified RIPA buffer,

containing 150 mM sodium chloride, 50 mM Tris HCl, pH 7.4, 1 mM EDTA, 1.0% NP-40, 0.5% sodium deoxycholic acid, 0.1% SDS and 0.01% (w/v) sodium azide to lyse the cells. Protein integrity was ensured using a cocktail of protease inhibitors with broad

specificity for the inhibition of aspartic, cysteine, and serine proteases as well

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
br>27709. United States

 ${\sf Email: info@biorbyt.com \mid Phone: +1 (415) \ 906-5211 \mid Fax: +1 \ (415) \ 651-8558}$