

Product Datasheet

HeLa Cell Nuclear Extract Etoposide Stimulated (orb348685)

Description

HeLa Cell Nuclear Extract Etoposide

Conjugation

Unconjugated

Tested

SDS-PAGE, WB

Applications

Preservatives

Preservative: None. Stabilizer: 10% (v/v) Glycerol. 1X SDS-PAGE Sample Buffer (62.5 mM Tris HCl, 2% SDS, 10% Glycerol and 0.005% bromophenol blue, pH 6.8)

Form/Appearance

Liquid (sterile filtered)

Concentration

1.0 mg/ml

Storage

Store HeLa Cell Nuclear Extract Etoposide Stimulated at -70° C or COLDER. For extended storage, aliquot Nuclear Extract to minimize freeze/thaw cycles.

Note

For research use only

Application notes

ready-to-use nuclear extracts are especially prepared as positive controls for separation by SDS-PAGE and subsequent western blot analysis. Nuclear extracts are supplied in denaturing buffer without dissociating agents. Heat nuclear extract to 95°C for 5 minutes and rapidly cool. If dissociating conditions are desired add reducing agent prior to heating. The recommended loading volume per lane is 10-30 ul depending on the size format of your gel.

Purity

The cells were grown in Eagles medium supplemented with 10% FBS (Fetal Bovine Serum). Cells were treated with 10 µg/ml Etoside for 13 h. The lysate was prepared by first washing the cells in PBS. Washed cells were then incubated on ice in lysis buffer containing 10 mM HEPES, 60 mM KCl, 1.0 mM EDTA, 0.075% (v/v) NP40 and 1.0 mM DTT, pH 7.6. Protein integrity is ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of serine, cysteine, and aspartic

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709, United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

collected and washed in lysis buffer minus detergent. Nuclei were lysed