

Product Datasheet

HeLa Cell Nuclear Extract TNFa Stimulated (orb348688)

Description

HeLa Cell Nuclear Extract TNFα

Conjugation	Unconjugated
Tested Applications	WB
Preservatives	Preservative: None. Stabilizer: 10% (v/v) Glycerol. 1X SDS-PAGE Sample Buffer (62.5 mM Tris HCl, 2% SDS, 10% Glycerol and 0.005% bromophenol blue, pH 6.8)
Form/Appearance	Liquid (sterile filtered)
Storage	Store HeLa Cell Nuclear Extract TNFα Stimulated at -70° C or COLDER. For extended storage, aliquot Nuclear Extract to minimize freeze/thaw cycles.
Note	For research use only
Application notes	ready-to-use nuclear extracts are especially prepared as positive controls for separation by SDS-PAGE and subsequent western blot analysis. Nuclear extracts are supplied in denaturing buffer without dissociating agents. Heat nuclear extract to 95°C for 5 minutes and rapidly cool. If dissociating conditions are desired add reducing agent prior to heating. The recommended loading volume per lane is 10-30 l depending on the size format of your gel.
Purity	The cells were grown in DMEM supplemented with 10% FBS (Fetal Bovine Serum). Cells were treated with 0.2 µg/ml TNFα for 30 min. The lysate was prepared by first washing the cells in PBS. Washed cells were then incubated on ice in lysis buffer containing 10 mM HEPES, 60 mM KCl, 1.0 mM EDTA, 0.075% (v/v) NP40 and 1.0 mM DTT, pH 7.6. Protein integrity is ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and serine proteases as well as aminopeptidases (0.1 mM AEBSF HCl, 0.08 µM Aprotinin, 5 µM Bestatin, 1.5

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709. United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

MgCl₂, 0.42 M NaCl, 0.2 mM EDTA,