

## Product Datasheet

### MOLT-4 Cell Nuclear Extract (orb348697)

## Description

MOLT-4 Cell Nuclear Extract

### Conjugation

Unconjugated

### Tested

ChIP, IP, SDS-PAGE, WB

### Applications

### Preservatives

Preservative: None. Stabilizer: None. 1X RIPA Buffer with HALT Protease and Phosphatase Inhibitors

### Form/Appearance

Liquid (sterile filtered)

### Concentration

4.0 mg/ml

### Storage

Store vial at -70° C or COLDER. For extended storage, aliquot Nuclear Extract to minimize freeze/thaw cycles.

### Note

For research use only

### Application notes

orb348697 has been tested by SDS-PAGE and western blot. Multi-purpose MOLT-4 nuclear extracts are especially prepared as positive control for multiple assays including western blot, immunoprecipitation (IP), capture ELISA or other assays requiring native protein sample. For separation by SDS-PAGE and subsequent western blot analysis, lysates should be diluted by user to desired concentration in SDS-PAGE buffer with 2-mercaptoethanol or dithiothreitol as the reducing agent and heated to 95° C for 5 minutes. Sample is ready for use in immunoprecipitation and ELISA experiments, conditions should be optimized by the user. Rockland recommends its TrueBlot IP reagents for immunoprecipitation experiments.

### Purity

MOLT-4 cells were grown in RPMI-1640 medium supplemented with 10% fetal bovine serum. Cells were washed with PBS and then incubated on ice in modified RIPA buffer, containing 150 mM sodium chloride, 50 mM Tris HCl, pH 7.4, 1 mM EDTA, 1.0% NP-40, 0.5% sodium deoxycholic acid, 0.1% SDS and 0.01% (w/v) sodium azide to lyse the cells. Protein integrity was ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and serine proteases as well as aminopeptidases (0.1 mM AEBSF, 0.02 mM Acetatin, 5 mM Bestatin, 1 mM

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive<br>Research Triangle Park<br>Durham, North Carolina<br>27709, United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com) | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

modified Lowry assay using a commercially available kit. Protein concentration was