

## **Product Datasheet**

HT-1080 Whole Cell Lysate (orb348706)



## www.biorbyt.com

Description hts. HT-1080 Whole Cell Lysate

**Conjugation** Unconjugated

Tested
Applications

ChIP, FC, IP, SDS-PAGE, WB

**Preservatives** Preservative: None. Stabilizer: None. 1X

RIPA Buffer with HALT Protease and

**Phosphatase Inhibitors** 

Form/Appearance Liquid (sterile filtered)

**Concentration** 4.0 mg/ml

**Storage** Store vial at -70° C or COLDER. For

extended storage, aliquot lysate to minimize freeze/thaw cycles.

**Note** For research use only

**Application notes** orb348706 has been tested by SDS-PAGE.

Multi-purpose HT-1080 Whole Cell Lysate are especially prepared as positive control for multiple assays including western blot, immunoprecipitation (IP), capture ELISA or other assays requiring native protein sample. For separation by SDS-PAGE and subsequent western blot analysis, lysates should be diluted by user to desired concentration in SDS-PAGE buffer with 2mercaptoethanol or dithiothreitol as the reducing agent and heated to 95° C for 5 minutes. Sample is ready for use in immunoprecipitation and ELISA experiments, conditions should be optimized by the user. Rockland recommends its TrueBlot IP reagents for

immunoprecipitation experiments.

**Purity** HT-1080 cells were grown in Eagle's

Minimum Essential Medium supplemented with 10% fetal bovine serum. Cells were washed with PBS and then incubated on ice in modified RIPA buffer, containing 150 mM sodium chloride, 50 mM Tris HCl, pH 7.4, 1 mM EDTA, 1.0% NP-40, 0.5% sodium deoxycholic acid , 0.1% SDS and 0.01% (w/v) sodium azide to lyse the cells. Protein integrity was ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and

serine proteases as well as

aminopeptidases (0.1 mM AEBSF HCI, 0.08

Biorbyt Ltd.

Biorbyt LLC.