

Product Datasheet

Mouse Embryonic Fibroblast Whole Cell Lysate (orb348722)

Description

Mouse Embryonic Fibroblast Whole

Conjugation

Unconjugated

Tested

SDS-PAGE, WB

Applications

Preservatives

Preservative: None. Stabilizer: 10% (v/v) Glycerol. 1X SDS-PAGE Sample Buffer (62.5 mM Tris HCl, 2% SDS, 10% Glycerol and 0.005% bromophenol blue, pH 6.8)

Form/Appearance

Liquid (sterile filtered)

Concentration

1.0 mg/ml

Storage

Store vial at -70° C or COLDER. For extended storage, aliquot contents to minimize freeze/thaw cycles.

Note

For research use only

Application notes

ready-to-use lysates are especially prepared as positive controls for separation by SDS-PAGE and subsequent western blot analysis. Lysates are prepared in denaturing buffer WITHOUT dissociating agents (i.e. no 2-mercaptoethanol or dithiothreitol has been added). Heat lysate to 95°C for 5 minutes and rapidly cool. If dissociating conditions are desired, add reducing agent prior to heating. The recommended loading volume per lane is 10-20 µl depending on the size format of your gel.

Purity

The cells were grown in RPMI supplemented with 10% FBS (Fetal Bovine Serum). The lysate was prepared by first washing the cells in PBS. Washed cells are then incubated on ice in modified RIPA buffer containing 150 mM sodium chloride, 50 mM Tris Cl, pH 7.4, 1 mM EDTA, 1.0% NP-40, 0.25% sodium deoxycholic acid to lyse the cells. Protein integrity is ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and serine

proteases as well as aminopeptidases

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280 240

Biorbyt LLC.

68 TW Alexander Drive
Research Triangle Park
Durham, North Carolina
27709, United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

1mM Na3VO4. Cell debris was removed by membrane filtration